

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C12N 15/11, 15/12, C07K 14/46, 19/00, 16/18, G01N 33/68, C12Q 1/68		A1	(11) International Publication Number: WO 99/15647 (43) International Publication Date: 1 April 1999 (01.04.99)
(21) International Application Number: PCT/AU98/00795 (22) International Filing Date: 23 September 1998 (23.09.98) (30) Priority Data: PO 9388 23 September 1997 (23.09.97) AU (71) Applicant (for all designated States except US): GARVAN INSTITUTE OF MEDICAL RESEARCH [AU/AU]; c/o St. Vincent's Hospital, 384 Victoria Street, Darlinghurst, NSW 2010 (AU). (72) Inventors; and (75) Inventors/Applicants (for US only): DALY, Roger, John [GB/AU]; 49 Gerard Street, Alexandria, NSW 2015 (AU). SUTHERLAND, Robert, Lyndsay [AU/AU]; 20 Northcote Road, Lindfield, NSW 2070 (AU). (74) Agent: F.B. RICE & CO.; 605 Darling Street, Balmain, NSW 2041 (AU).			(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published With international search report.
(54) Title: A POTENTIAL EFFECTOR FOR THE GRB7 FAMILY OF SIGNALLING PROTEINS			
(57) Abstract A novel polynucleotide molecule is disclosed which encodes a candidate effector protein for the Grb7 family of signalling proteins. Detection of the protein in a sample such as a homogenised tissue sample should provide a useful tumour marker and/or prognostic indicator for certain human cancers such as breast and prostate cancer.			

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

A POTENTIAL EFFECTOR FOR THE GRB7 FAMILY OF SIGNALLING PROTEINS

Field of the Invention:

5

The present invention relates to a novel polynucleotide molecule encoding a candidate effector protein for the Grb7 family of signalling proteins. Detection of the encoded protein in a tissue sample should provide a useful tumour marker and/or prognostic indicator. Furthermore, antagonism of the interaction between Grb7 family members and the encoded protein should provide a novel treatment strategy for human diseases exhibiting aberrant receptor tyrosine kinase (RTK) signalling (e.g. cancer).

Background of the Invention

RTKs play a major role in the regulation of cellular growth, differentiation, motility and metabolism by converting an extracellular signal in the form of the binding of a specific hormone or growth factor to the activation of specific signalling pathways and hence modes of intracellular communication (Schlessinger and Ullrich, *Neuron* 9. 383-391, 1992). Activation of RTKs results in both autophosphorylation of the receptor and the phosphorylation of downstream targets on tyrosine residues. It has become evident over the last decade that key elements in receptor-substrate and other protein-protein interactions in RTK signalling are src homology (SH)2 domains. SH2 domains are conserved modules of approximately 100 amino acids found in a wide variety of signalling molecules which bind to short tyrosine-phosphorylated peptide sequences. The specificity of interaction is determined both by the nature of the amino acids flanking the phosphotyrosine residue in the target peptide and residues in the SH2 domain which interact with these sites (Pawson, *Nature* 373, 573-580, 1995).

SH2-domain containing proteins can be divided into two classes: those which possess a catalytic function (e.g. the cytoplasmic tyrosine kinase c-src and the tyrosine phosphatase SH-PTP2) and those which consist entirely of non-catalytic protein domains (eg Grb2), the adaptor sub-class. The function of the latter class is to link separate catalytic subunits to a tyrosine-

phosphorylated receptor or signalling intermediate, and other non-catalytic protein modules are often involved in these interactions. For example, SH3 and WW domains (conserved regions of approximately 50 and 40 amino acids, respectively) bind proline-rich peptide ligands, and pleckstrin
5 homology domains (approximately 100 amino acids) interact with both specific phospholipid and protein targets (Pawson, 1995 *supra*).

The Grb7 family represents a family of SH2 domain-containing adaptors which currently contains three members: Grb7, 10 and 14 (Margolis
10 *et al*, *Proc. Natl. Acad. Sci. USA* 89, 8894-8898, 1992; Stein *et al*, *EMBO J* 13, 1331-1340, 1994; Ooi *et al*, *Oncogene* 10, 1621-1630, 1995; Daly *et al*, *J. Biol. Chem.* 271, 12502-12510, 1996). These proteins share a common overall architecture, consisting of an N-terminal region containing a highly conserved proline-rich decapeptide motif, a central region harbouring a PH domain and a C-terminal SH2 domain. The central region of approximately
15 300 amino acids bears significant homology to the *C. elegans* protein mig10, which is required for long range neuronal migration in embryos, otherwise the Grb7 family and mig10 are structurally distinct. However, they exhibit differences in both SH2 selectivity towards RTKs (Janes *et al*, *J. Biol. Chem.* 272, 8490-8497, 1997) and tissue distribution. The family has therefore
20 evolved to link particular receptors to downstream effectors in a tissue-specific manner. Interestingly, the genes encoding this family appear to have co-segregated with *ERBB* family genes during evolution. Thus *GRB7*, 10 and 14 are linked to *ERBB2*, *ERBB1* (epidermal growth factor receptor) and *ERBB4*, respectively (Stein *et al* 1994 *supra*; Ooi *et al*, 1995 *supra*; Baker *et al*,
25 *Genomics* 36, 218-220, 1996). The juxtaposition of *GRB7* and *ERBB2* leads to common co-amplification in human breast cancers, and since the two gene products are functionally linked, likely up-regulation of an undefined *erbB2* signalling pathway. Furthermore, *GRB14* also exhibits differential expression in human breast cancers (Daly *et al*, 1996 *supra*). These two proteins may
30 therefore modulate RTK signalling in this disease.

In order to identify proteins which bind to this family and therefore identify candidate effectors, we performed a genetic screen using the yeast two hybrid system and Grb14 "bait". This application describes the cloning and characterization of a novel interacting protein, currently designated
35 2.2412.

Disclosure of the Invention:

Thus, in a first aspect, the present invention provides an isolated polynucleotide molecule encoding a candidate effector protein for the Grb7 family of signalling proteins, wherein the polynucleotide molecule comprises
5 a nucleotide sequence having at least 75% sequence identity to that shown as SEQ ID NO: 1.

Preferably, the polynucleotide molecule comprises a nucleotide sequence having at least 85%, more preferably at least 95%, sequence
10 identity to that shown as SEQ ID NO: 1. Most preferably, the polynucleotide molecule comprises a nucleotide sequence encoding a polypeptide comprising an amino acid sequence substantially corresponding to that shown as SEQ ID NO: 2.

In a preferred embodiment of the invention of the first aspect, the
15 polynucleotide molecule comprises a nucleotide sequence which substantially corresponds to that shown as SEQ ID NO: 1.

The polynucleotide molecule may be a dominant negative mutant which encodes a gene product causing an altered phenotype by, for example, reducing or eliminating the activity of endogenous effector proteins of the
20 Grb7 family of signalling proteins.

The polynucleotide molecule may be incorporated into plasmids or expression vectors (including viral vectors), which may then be introduced into suitable host cells such as bacterial, yeast, insect and mammalian host cells. Such host cells may be used to express the protein encoded by the
25 polynucleotide molecule.

Accordingly, in a second aspect, the present invention provides a host cell transformed with the polynucleotide molecule of the first aspect.

In a third aspect, the present invention provides a method of producing a protein, comprising culturing the host cell of the second aspect under
30 conditions suitable for the expression of the polynucleotide molecule and optionally recovering the protein.

Preferably, the host cell is mammalian or of insect origin. Where the cell is mammalian, it is presently preferred that it be a Chinese hamster ovary (CHO) cell or human embryonic kidney (HEK) 293 cell. Where the
35 host cell is of insect origin, it is presently preferred that it be an insect Sf9 cell.

In a fourth aspect, the present invention provides a purified protein encoded by the polynucleotide molecule of the first aspect.

In a preferred embodiment of this aspect, the purified protein comprises an amino acid sequence substantially corresponding to that shown
5 as SEQ ID NO: 2.

In a fifth aspect, the present invention provides a fusion protein comprising an amino acid sequence substantially corresponding to that shown as SEQ ID NO: 2.

Fusion proteins according to the fifth aspect may include an N-
10 terminal fragment of a protein such as β -galactosidase to assist in the expression and selection of host cells expressing candidate effector protein, or may include a functional fragment of any other suitable protein to confer additional activity(ies).

In a sixth aspect, the present invention provides an antibody or
15 fragment thereof which specifically binds to the protein of the fourth aspect.

The antibody may be monoclonal or polyclonal, however, it is presently preferred that the antibody is a monoclonal antibody. Suitable antibody fragments include Fab, F(ab')₂ and scFv.

In a seventh aspect, the present invention provides an oligonucleotide
20 probe comprising a nucleotide sequence of at least 12 nucleotides, the oligonucleotide probe comprising a nucleotide sequence such that the oligonucleotide probe selectively hybridises to the polynucleotide molecule of the first aspect under high stringency conditions (Sambrook *et al.*, *Molecular Cloning: a Laboratory Manual*, Second Edition, Cold Spring Harbor
25 Laboratory Press).

In a preferred embodiment of this aspect, the oligonucleotide probe is labelled. In a further preferred embodiment of this aspect, the oligonucleotide probe comprises a nucleotide sequence of at least 18 nucleotides.

30 In an eighth aspect, the present invention provides a method of detecting in a sample the presence of an effector protein for the Grb7 family of proteins, the method comprising reacting the sample with an antibody or fragment thereof of the sixth aspect, and detecting the binding of the antibody or fragment thereof.

The method of the eighth aspect may be conducted using any immunoassays well known in the art (e.g. ELISA). The sample may be, for example, a cell lysate or homogenate prepared from a tissue biopsy.

In a ninth aspect, the present invention provides a method of detecting
5 in a sample the presence of mRNA encoding an effector protein for the Grb7 family of proteins, the method comprising reacting the sample with an oligonucleotide probe of the seventh aspect, and detecting the binding of the probe.

The method of the ninth aspect may be conducted using any
10 hybridisation assays well known in the art (e.g. Northern blot). The sample may be a poly(A) RNA preparation or homogenate prepared from a tissue biopsy.

Grb7 family proteins exhibit differential expression in certain human cancers (particularly breast and prostate cancer) and may therefore be
15 involved in tumour progression. Detection of the protein encoded by the cDNA 2.2412 in a sample should provide a useful tumour marker and/or prognostic indicator for these cancers. Furthermore, the interaction of Grb7 family members with 2.2412 may provide a novel target for therapeutic intervention.

20 It is to be understood that methods of detecting suitable agonists and methods of therapy utilising detected agonists also form part of the present invention.

The term "substantially corresponds" as used herein in relation to the nucleotide sequence shown as SEQ ID NO: 1 is intended to encompass minor
25 variations in the nucleotide sequence which due to degeneracy in the DNA code do not result in a change in the encoded protein. Further, this term is intended to encompass other minor variations in the sequence which may be required to enhance expression in a particular system but in which the variations do not result in a decrease in biological activity of the encoded
30 protein.

The term "substantially corresponding" as used herein in relation to the amino acid sequences shown as SEQ ID NO: 2 is intended to encompass minor variations in the amino acid sequences which do not result in a decrease in biological activity of the protein. These variations may include
35 conservative amino acid substitutions. The substitutions envisaged are:-

G, A, V, I, L, M: D, E; N, Q; S, T; K, R, H; F, Y, W, H; and

P, N α -alkalamino acids.

The terms "comprise", "comprises" and "comprising" as used throughout the specification are intended to refer to the inclusion of a stated step, component or feature of group of steps, components of features with or without the inclusion of a further step, component or feature or group of steps, components or features.

The invention will hereinafter be described with reference to the accompanying figure and the following, non-limiting example.

Brief description of the accompanying figure:

Figure 1 provides the nucleotide and amino acid (single letter code) sequence of 2.2412. Numbers refer to distances in base pairs. Ankyrin-type repeat sequences are underlined. An additional repeat sequence is indicated by italics.

The stop codon is represented by an asterisk. The original cDNA clone 2.2412 isolated by the two hybrid screen spans nucleotides 694-2664 of this sequence.

Figure 2 provides a map of the 2.2412-binding region on Grb14.

A. Structure of the deletion constructs used in the analysis. Gal4 DNA-BD fusion constructs encoding full length Grb14 (FL), the N-terminal (N), central region (C) and N-terminal + central region (N + C) were generated in the vector pAS2.1.

B. Results of β -galactosidase activity assays following transformation of the above plasmids into yeast strain Y190 together with the original 2.2412 cDNA clone in pACT-2.

Example: CLONING AND CHARACTERISATION OF 2.2412

Yeast two hybrid screen

The yeast two hybrid system exploits protein-protein interactions to reconstitute a functional transcriptional activator which can then be detected using a gene reporter system (Fields and Sternglanz, *TIG*, 10, 286-292, 1994). The technique takes advantage of the properties of the Gal4 protein of the yeast *S. cerevisiae*. The Gal4 DNA binding domain (DNA-BD) or activation domain (AD) alone are incapable of inducing transcription. However, an interaction between two proteins synthesized as DNA-BD- and AD-fusions, respectively, brings the Gal4 domains into close proximity and results in

transcriptional activation of two reporter genes (*HIS3* and *LacZ*) which can be monitored by growth on selective medium and biochemical assays.

A plasmid construct encoding a Gal4 DNA-BD-Grb14 fusion was generated as follows. The plasmid *GRB14/pRcCMVF* containing full length *GRB14* cDNA (Daly *et al.*, 1996) was restricted with *HindIII* and Klenow treated to create blunt ends, and then digested with *BclI* to release three fragments of approximately 1.1, 4.2 and 1.7 kb. The 1.7 kb fragment was isolated and cloned into the *NdeI* (Klenow treated) and *BamHI* sites of the yeast expression vector pAS2.1 (Clontech) to generate *GRB14/pAS2.1* containing an in-frame fusion of full length Grb14 with the GAL4 DNA-BD. This construct was introduced by electroporation into the yeast strain CG1945 (*MATa*, *ura3-52*, *his3-200*, *ade2-101*, *lys2-801*, *trp1-901*, *leu2-3, 112*, *gal4-542*, *gal80-538*, *cyh^r2*, *LYS2::GAL1UAS-GAL1TATA-HIS3*, *URA3::GAL417mers(x3)-CYC1TATA-lacZ*) selecting for tryptophan prototrophy. The expression of the fusion protein was verified by Western blot analysis with antibodies directed against the Flag epitope and the Gal4 DNA-BD. The recipient strain was then grown to mid-log phase and a human liver cDNA library in the vector pACT2 (Clontech) introduced using the LiAc procedure (Schiestl and Gietz, *Curr. Genet.* 16, 339-346, 1989). Transformants were then selected for tryptophan, leucine and histidine prototrophy in the presence of 5mM 3-aminotriazole.

From a screen of 1×10^6 clones, 39 colonies were initially selected on synthetic complete (SC)-leu-his-trp + 3AT medium and were then tested for β -galactosidase activity. 12 clones scored positive in the latter assay and were subjected to cycloheximide (CHX) curing to remove the bait plasmid by streaking out on SC-leu media containing 10ug/ml CHX (pAS2-1 contains the *CYH2* gene which restores CHX sensitivity to CG1945 cells). This enabled confirmation of the bait dependency of *LacZ* activation and subsequent isolation of the pACT2 plasmids encoding interacting proteins by standard methodology (Philippsen *et al.*, *Methods in Enzymology* 194, 170-177). Back transformations were then performed in which these pACT2 plasmids were introduced into CG1945 strains containing the bait plasmid (*GRB14/pAS2-1*) or constructs encoding non-related Gal4 DNA-BD fusions in order to confirm the specificity of the interactions.

The DNA sequences of the cDNA inserts were then obtained by cycle sequencing (f-mol kit, Promega) using pACT2-specific and/or clone-specific

primers. Based on their nucleotide sequences the 12 interacting clones were classified into 6 independent groups (see Table I).

TABLE I: Characterization of cDNA clones isolated by the yeast two hybrid screen.

	Class	No. of clones	Identity	Mean RLU (Liquid assay)	Colour intensity (Filter assay)
10	1	6	Nedd4	2.86×10^6	++++
	2	2	Htk	1.86×10^5	++
	3	1	2.2412	5.18×10^6	++++
	4	1	Proleosome	3.88×10^2	+/-
	5	1	Somatostatin	1.45×10^3	+/-
15			receptor		
	6	1	L-arginine:glycine amidinotransferase	8.61×10^2	+/-

The 12 clones exhibiting activation of both the *HIS3* and *lacZ* reporter genes were divided into 6 groups by sequence analysis of their cDNA inserts. Results of β -galactosidase activity assays performed using two methodologies are shown. The liquid culture-derived method (Galacto-Light, TROPIX) is more quantitative: results are given in mean relative light units (RLU) and are normalized for the protein content of the samples. Blue/white screening of the cDNA clones was also performed using a colony lift filter assay (Clontech). The intensity of blue colour development over approximately 2h is scored from +/- (very weak) to ++++ (strong).

Six clones were partial cDNAs corresponding to Nedd4, a multidomain protein containing a calcium-dependent phospholipid binding (CaLB) domain, four WW domains and a C-terminal region homologous to the E6-AP carboxyl-terminus (Kumar *et al*, *Biochem. Biophys. Res. Commun.* 185, 1155-1161, 1992; Sudol *et al* *J. Biol. Chem.* 270, 14733-14741, 1995; Huibregtse *et al* *Proc. Natl. Acad. Sci. USA* 92, 2563-2567, 1995). The latter is likely to confer E3 ubiquitin-protein ligase activity on Nedd4. The pACT2 clones isolated encoded the CaLB domain together with the first 22 amino acids of the first WW domain.

Two clones encoded the intracellular region and part of the extracellular domain of Htk, which is a RTK of the Eph family (Bennett *et al* *J. Biol. Chem.* 269, 14211-14218, 1994). The recruitment of Grb14 by Htk is of interest for two reasons. First, the expression profile of both Htk and the murine homologue myk-1 are indicative of a potential role in mammary gland development and neoplasia (Andres *et al* *Oncogene* 9, 1461-1467, 1994; Berclaz *et al* *Biochem. Biophys. Res. Comm.* 226, 869-875, 1996). Second, Eph family members may be involved in the regulation of cell migration (Tessier-Lavigne, *Cell* 82, 345-348, 1995), which is intriguing given the homology of the Grb7 family to the *C. elegans* protein mig10 (Stein *et al.* 1994 *supra*).

A novel cDNA of 1971 bp, designated 2.2412, was also isolated. This clone encoded a polypeptide of 657 amino acids in frame with the Gal4 DNA-BD. The cDNA did not contain a stop codon, and this, together with the Northern analysis described below, indicated that it was incomplete. This DNA fragment was therefore used as a probe to screen a human placental cDNA library (5' STRETCH PLUS, Clontech, in λ gt10). This resulted in the isolation of two clones, designated clone 8 and clone 12. Clone 8 was approximately 2 kb and overlapped the original 2.2412 clone by 900 bp at the 3' end. This clone provided the carboxy-terminal end of the 2.2412 protein sequence (Figure 1). Clone 12 was approximately 3.5 kb and to date has provided an additional 692 bp of sequence information in the 5' direction. The nucleotide and protein sequence for 2.2412 provided by these overlapping clones is shown in Figure 1. Since a 5' initiation codon has yet to be identified the coding sequence still appears to be incomplete.

Further characterization of 2.2412

Database searches using the 2.2412 cDNA sequence revealed significant homology with a large number of proteins containing ankyrin-like repeats. These sequences were first identified as homologous regions between certain cell cycle regulatory proteins and the *Drosophila* protein Notch (Breedon and Nasmyth, *Nature* 329, 651-654, 1987) but subsequently they have been identified in a wide variety of other proteins where they are thought to function in protein-protein interactions (Bork, *Proteins* 17, 363-374, 1993). Subsequent analysis of the protein sequence identified 18 consecutive ankyrin repeats and an additional repetitive element (Figure 1).

The ankyrin repeat region is followed by a stretch of approximately 40 amino

acids rich in serine residues. The remaining C-terminal region has a relatively high content of charged amino acids.

Northern analysis of 2.2412 mRNA expression

5 Northern blot analysis of multiple tissue northern (Clontech) was performed using the original 2.2412 cDNA as a probe. This resulted in the detection of a single mRNA transcript of approximately 7 kb in all tissues examined with the exception of the kidney. Expression was particularly high in skeletal muscle and placenta. The size of this transcript compared to that
10 of the 2.2412 clone indicates that the latter represents only a partial cDNA.

Genomic localization of the 2.2412 gene

Fluorescence *in situ* hybridization of the original 2.2412 cDNA to normal metaphases (Baker *et al*, 1996 *supra*) and reference to the FRA10A
15 fragile site at 10q23.32 localized the gene to between chromosome 10q23.2 and proximal 10q23.32. Interestingly, deletions in the 10q22-25 region of chromosome 10 have been detected in a variety of human cancers including breast, prostate, renal, small cell lung and endometrial carcinomas, glioblastoma multiforme, melanoma and meningiomas, suggesting the
20 presence of one or more tumour suppressive loci in this region (Li *et al*, *Science* 275, 1943-1947, 1997; Steck *et al*, *Nature Genetics* 15, 356-362, 1997, and references therein). Two candidate tumour suppressor genes have been identified in this region (MMAC1/PTEN and MXI1. Li *et al* 1997 *supra*; Steck *et al* 1997 *supra*; Albarosa *et al*, *Hum. Genet.* 95, 709-711, 1995).

25

Analysis of the interaction between 2.2412 and Grb7 family members

cDNAs encoding the full length and N- and C-terminal regions of the original 2.2412 cDNA clone (nucleotides 694-2664, 694-1614 and 1615-2664 of the sequence shown in Figure 1, respectively) were cloned into the vector
30 pGEX4T2 (Pharmacia). The full length construct was generated by subcloning from the pACT2 clone as a NdeI fragment, whereas the shorter constructs were synthesized by directional cloning of PCR products. The corresponding GST-fusion proteins were purified from IPTG-induced bacterial cultures using glutathione-agarose beads (Smith and Johnson, *Gene*
35 67, 31-40, 1988). These immobilized fusion proteins were then incubated with lysates from cells expressing Flag epitope-tagged Grb14 (Daly *et al*, 1996

supra) or human breast cancer cells expressing high levels of Grb7 (SK-BR-3: Stein *et al.*, 1994) as described previously (Daly *et al.*, 1996). Following washing, bound proteins were detected by Western blot analysis. The results indicated that 2.2412 bound specifically to both Grb14 and Grb7 *in vitro*, and
5 that the N-terminal fusion protein bound more strongly than that derived from the C-terminus. These data, obtained using a different methodology for detecting protein-protein interactions to the yeast two hybrid system, confirm that 2.2412 interacts with Grb14. Furthermore, 2.2412 also binds Grb7. Consequently 2.2412 appears to represent a general effector for the Grb7
10 family.

Mapping of the 2.2412 binding region on Grb14

In order to identify the region of Grb14 that interacts with 2.2412, a series of Grb14 deletion mutants were generated by cloning PCR fragments
15 synthesized using the appropriate flanking primers into the vector pAS2.1. These fragments spanned the following regions: N-terminus ("N", amino acids 1-110), the central region ("C") encompassing the mig10 homology and the "between PH and SH2" (BPS) domain (amino acids 110-437) and the N-terminal and central regions ("N + C", amino acids 1-437). These plasmids
20 were individually transformed into the yeast strain Y190 (*MATa*, *ura3-52*, *his3-200*, *ade2-101*, *lys2-801*, *trp1-901*, *leu2-3, 112*, *gal4Δ*, *gal80Δ*, *cyh^r2*, *LYS2::GAL1UAS-HIS3TATA-HIS3*, *URA3::GAL1UAS-GAL1TATA-lacZ*) and expression of the appropriately sized Gal4 DNA-BD fusion proteins confirmed by Western blotting. Following transformation of the resulting
25 yeast strains with the original 2.2412 cDNA clone in pACT-2, the strength of the interaction was determined by either liquid- or filter-based β -galactosidase assays. The results are presented in Figure 2, and demonstrate that the N-terminal region of Grb14 is not only required, but is also sufficient, for binding 2.2412. This supports the hypothesis that 2.2412
30 represents a general effector for the Grb7 family, since the N-terminal region of these proteins contains a highly conserved proline-rich motif which may mediate this interaction.

It will be appreciated by persons skilled in the art that numerous variations and/or modifications may be made to the invention as shown in the specific embodiments without departing from the spirit or scope of the invention as broadly described. The present embodiments are, therefore, to
5 be considered in all respects as illustrative and not restrictive.

Sequence listings:

SEQUENCE LISTING

Applicant: Garvan Institute of Medical Research

Title of Invention: A potential effector for the Grb7 family of signalling proteins.

Current Application Number:

Current Filing Date:

Prior Application Number: PO9388

Prior Application Filing Date: 1997-09-23

Number of ID SEQ Nos: 2

Software: PatentIn Ver. 2.0

SEQ ID NO: 1

Length: 3400

Type: DNA

Organism: Homo sapiens

Sequence: 1

```

attcctcttc ataatgcatg ctcttttggg catgctgaag tagtcaatct cctttttgcga 60
catggtgcag accccaatgc tcgagataat tggaattata ctctctcca tgaagctgca 120
attaaaggaa agattgatgt ttgcattgtg ctgttacagc atggagctga gccaaccatc 180
cgaaatacag atggaaggac agcattggat ttagcagatc catctgccaa agcagtgcct 240
actggtgaat ataagaaaaga tgaactctta gaaagtgccg ggagtggcaa tgaagaaaaa 300
atgatggctc tactcacacc attaaatgtc aactgccacg caagtgatgg cagaaagtca 360
actccattac atttggcagc aggatataac agagtaaaga ttgtacagct gttactgcaa 420
catggacgtg atgtccatgc taaagataaa ggtgatctgg taccattaca caatgcctgt 480
tcttatggtc attatgaagt aactgaactt ttggtcaagc atgggtggctg tgtaaatgca 540
atggacttgt ggcaattcac tcctcttcac gaggcagctt ctaagaacag ggttgaagta 600
tggtctcttc tcttaagtta tgggtgcagac ccaacactgc tcaattgtaa gaataaaagt 660
gctatagact tggctcccac accacagtta aaagaaagat tagcatatga atttaaaggc 720
cactcggtgc tgcaagctgc acgagaagct gatgttactc gaatcaaaaa acatctctct 780
ctggaaatgg tgaatttcaa gcacctcaa acacatgaaa cagcattgca ttgtgctgct 840
gcacctccat atcccaaaag aaagcaaata tgtgaactgt tgctaagaaa aggagcaaac 900
atcaatgaaa agactaaaga attcttgact cctctgcacg tggcatctga gaaagctcat 960
aatgatgttg ttgaagtagt ggtgaaacat gaagcaaagg ttaatgctct ggataatctt 1020
ggtcagactt ctctacacag agctgcatat tgtggtcatc tacaaacctg ccgcctactc 1080
ctgagctatg ggtgtgatcc taacattata tcccttcagg gctttactgc tttacagatg 1140
ggaaatgaaa atgtacagca actcctccaa gagggatatc cattaggtaa ttcagaggca 1200
gacagacaat tgctggaagc tgcaaaggct ggagatgtcg aaactgtaaa aaaactgtgt 1260
actgttcaga gtgtcaactg cagagacatt gaagggcgct agtctacacc acttcatttt 1320
gcagctgggt ataacagagt gtccgtgggt gaatatctgc tacagcatgg agctgatgtg 1380
catgctaaag ataaaggagg ccttgtacct ttgcacaatg catgttctta cggacattat 1440
gaagttgcag aacttcttgt taaacatgga gcagtagtta atgtagctga tttatggaaa 1500
tttacacctt tactgaagc agcagcaaaa ggaaaatatg aaatttgcaa acttctgctc 1560
cagcatgggtg cagaccctac aaaaaaaaaa agggatggaa atactccttt ggatcttgtt 1620
aaagatggag atacagatat tcaagatctg cttaggggag atgcagcttt gctagatgct 1680
gccaagaagg gttgttttagc cagagtgaag aagttgtctt ctctgataa tgtaaattgc 1740
cgcgataccc aaggcagaca ttcaacacct ttacatttag cagctgggta taataattta 1800
gaagttgcag agtatttgtt acaacacgga gctgatgtga atgccaaga caaaggagga 1860
cttattcctt tacataatgc agcatcttac gggcatgtag atgtagcagc tctactaata 1920

```

```

aagtataatg catctctcaa tgccacggac aaatgggctt tcacaccttt gcacgaagca 1980
gccccaaagg gacgaacaca gctttgtgct ttgttgctag cccatggagc tgacccgact 2040
cttaaaaaatc aggaaggaca aacaccttta gatttagttt cagcagatga tgtcagcgct 2100
cttctgacag cagccatgcc cccatctgct ctgccctctt gttacaagcc tcaagtgtct 2160
aatgggtgtga gaagcccagg agccactgca gatgctctct cttcaggtcc atctagccca 2220
tcaagccttt ctgcagccag cagtcttgac aacttatctg ggagtttttc agaactgtct 2280
tcagtagtta gttcaagtgg aacagagggg gcttccagtt tggagaaaaa ggaggttcca 2340
ggagtagatt ttagcataac tcaattcgta aggaatcttg gacttgagca cctaattgat 2400
atatttgaga gagaacagat cactttggat gtattagtgt agatggggca caaggagctg 2460
aaggagattg gaatcaatgc ttatggacat aggcacaaac taattaaagg agtcgagaga 2520
cttatctccg gacaacaagg tcttaacca tatttaactt tgaacacctc tggtagtgga 2580
acaattctta tagatctgtc tcctgatgat aaagagtttc agtctgtgga ggaagagatg 2640
caaagtacag ttcgagagca cagagatgga ggtcatgcag gtggaatctt caacagatac 2700
aatattctca agattcagaa ggtttgtaac aagaaactat gggaaagata cactcaccgg 2760
agaaaagaag tttctgaaga aaaccacaac catgccaatg aacgaatgct atttcatggg 2820
tctccttttg tgaatgcaat tatccacaaa ggctttgatg aaaggcatgc gtacataggt 2880
ggtatgtttg gagctggcat ttattttgct gaaaactctt ccaaaagcaa tcaatatgta 2940
tatggaattg gaggaggtac tgggtgtcca gttcacaaag acagatcttg ttacatttgc 3000
cacaggcagc tgctcttttg cgggtaacc ttgggaaagt ctttcctgca gttcagtgca 3060
atgaaaatgg cacattctcc tccaggtcat cactcagtea ctggtaggcc cagtgtaaat 3120
ggcctagcat tagctgaata tgttatttac agaggagaac aggccttatc tgagtattta 3180
attacttacc agattatgag gcctgaaggt atggtcgatg gataaatagt tattttaaga 3240
aactaattcc actgaacctt aaatcatcaa agcagcagtg gcctctacgt tttactcctt 3300
tgctgaaaaa aaatcatctt gccacaggc ctgtggcaaa aggataaaaa tgtgaacgaa 3360
gtttaacatt ctgacttgat aaagctttta taatgtacag 3400

```

SEQ ID NO: 2

Length: 1074

Type: PRT

Organism: Homo sapiens

Sequence: 2

```

Ile Pro Leu His Asn Ala Cys Ser Phe Gly His Ala Glu Val Val Asn
  1                      5              10              15

```

```

Leu Leu Leu Arg His Gly Ala Asp Pro Asn Ala Arg Asp Asn Trp Asn
    20              25              30

```

```

Tyr Thr Pro Leu His Glu Ala Ala Ile Lys Gly Lys Ile Asp Val Cys
    35              40              45

```

```

Ile Val Leu Leu Gln His Gly Ala Glu Pro Thr Ile Arg Asn Thr Asp
    50              55              60

```

```

Gly Arg Thr Ala Leu Asp Leu Ala Asp Pro Ser Ala Lys Ala Val Leu
    65              70              75              80

```

```

Thr Gly Glu Tyr Lys Lys Asp Glu Leu Leu Glu Ser Ala Arg Ser Gly
    85              90              95

```

```

Asn Glu Glu Lys Met Met Ala Leu Leu Thr Pro Leu Asn Val Asn Cys
   100              105              110

```

```

His Ala Ser Asp Gly Arg Lys Ser Thr Pro Leu His Leu Ala Ala Gly
   115              120              125

```

```

Tyr Asn Arg Val Lys Ile Val Gln Leu Leu Leu Gln His Gly Arg Asp
   130              135              140

```

```

Val His Ala Lys Asp Lys Gly Asp Leu Val Pro Leu His Asn Ala Cys

```


15

145		150		155		160
Ser Tyr Gly His Tyr Glu Val Thr Glu Leu Leu Val Lys His Gly Gly						
	165			170		175
Cys Val Asn Ala Met Asp Leu Trp Gln Phe Thr Pro Leu His Glu Ala						
	180			185		190
Ala Ser Lys Asn Arg Val Glu Val Cys Ser Leu Leu Leu Ser Tyr Gly						
	195			200		205
Ala Asp Pro Thr Leu Leu Asn Cys Lys Asn Lys Ser Ala Ile Asp Leu						
	210			215		220
Ala Pro Thr Pro Gln Leu Lys Glu Arg Leu Ala Tyr Glu Phe Lys Gly						
	225			230		235
His Ser Leu Leu Gln Ala Ala Arg Glu Ala Asp Val Thr Arg Ile Lys						
	245			250		255
Lys His Leu Ser Leu Glu Met Val Asn Phe Lys His Pro Gln Thr His						
	260			265		270
Glu Thr Ala Leu His Cys Ala Ala Ala Ser Pro Tyr Pro Lys Arg Lys						
	275			280		285
Gln Ile Cys Glu Leu Leu Leu Arg Lys Gly Ala Asn Ile Asn Glu Lys						
	290			295		300
Thr Lys Glu Phe Leu Thr Pro Leu His Val Ala Ser Glu Lys Ala His						
	305			310		315
Asn Asp Val Val Glu Val Val Val Lys His Glu Ala Lys Val Asn Ala						
	325			330		335
Leu Asp Asn Leu Gly Gln Thr Ser Leu His Arg Ala Ala Tyr Cys Gly						
	340			345		350
His Leu Gln Thr Cys Arg Leu Leu Leu Ser Tyr Gly Cys Asp Pro Asn						
	355			360		365
Ile Ile Ser Leu Gln Gly Phe Thr Ala Leu Gln Met Gly Asn Glu Asn						
	370			375		380
Val Gln Gln Leu Leu Gln Glu Gly Ile Ser Leu Gly Asn Ser Glu Ala						
	385			390		395
Asp Arg Gln Leu Leu Glu Ala Ala Lys Ala Gly Asp Val Glu Thr Val						
	405			410		415
Lys Lys Leu Cys Thr Val Gln Ser Val Asn Cys Arg Asp Ile Glu Gly						
	420			425		430
Arg Gln Ser Thr Pro Leu His Phe Ala Ala Gly Tyr Asn Arg Val Ser						
	435			440		445
Val Val Glu Tyr Leu Leu Gln His Gly Ala Asp Val His Ala Lys Asp						
	450			455		460
Lys Gly Gly Leu Val Pro Leu His Asn Ala Cys Ser Tyr Gly His Tyr						

465		470		475		480
Glu Val Ala Glu	Leu Leu Val Lys His	Gly Ala Val Val Asn Val Ala				
	485	490			495	
Asp Leu Trp Lys	Phe Thr Pro Leu His	Glu Ala Ala Ala Lys Gly Lys				
	500	505			510	
Tyr Glu Ile Cys	Lys Leu Leu Leu Gln His	Gly Ala Asp Pro Thr Lys				
	515	520			525	
Lys Asn Arg Asp	Gly Asn Thr Pro Leu Asp	Leu Val Lys Asp Gly Asp				
	530	535			540	
Thr Asp Ile Gln	Asp Leu Leu Arg Gly Asp	Ala Ala Leu Leu Asp Ala				
	545	550			555	
Ala Lys Lys Gly	Cys Leu Ala Arg Val Lys	Lys Leu Ser Ser Pro Asp				
	565	570			575	
Asn Val Asn Cys	Arg Asp Thr Gln Gly	Arg His Ser Thr Pro Leu His				
	580	585			590	
Leu Ala Ala Gly	Tyr Asn Asn Leu Glu Val	Ala Glu Tyr Leu Leu Gln				
	595	600			605	
His Gly Ala Asp	Val Asn Ala Gln Asp Lys	Gly Gly Leu Ile Pro Leu				
	610	615			620	
His Asn Ala Ala	Ser Tyr Gly His Val Asp	Val Ala Ala Leu Leu Ile				
	625	630			635	
Lys Tyr Asn Ala	Ser Leu Asn Ala Thr Asp	Lys Trp Ala Phe Thr Pro				
	645	650			655	
Leu His Glu Ala	Ala Gln Lys Gly Arg Thr	Gln Leu Cys Ala Leu Leu				
	660	665			670	
Leu Ala His Gly	Ala Asp Pro Thr Leu Lys	Asn Gln Glu Gly Gln Thr				
	675	680			685	
Pro Leu Asp Leu	Val Ser Ala Asp Asp Val	Ser Ala Leu Leu Thr Ala				
	690	695			700	
Ala Met Pro Pro	Ser Ala Leu Pro Ser Cys	Tyr Lys Pro Gln Val Leu				
	705	710			715	
Asn Gly Val Arg	Ser Pro Gly Ala Thr Ala	Asp Ala Leu Ser Ser Gly				
	725	730			735	
Pro Ser Ser Pro	Ser Ser Leu Ser Ala Ala	Ser Ser Leu Asp Asn Leu				
	740	745			750	
Ser Gly Ser Phe	Ser Glu Leu Ser Ser Val	Val Ser Ser Ser Gly Thr				
	755	760			765	
Glu Gly Ala Ser	Ser Leu Glu Lys Lys Glu	Val Pro Gly Val Asp Phe				
	770	775			780	
Ser Ile Thr Gln	Phe Val Arg Asn Leu Gly	Leu Glu His Leu Met Asp				

785		790		795		800
Ile Phe Glu Arg	Glu Gln Ile Thr Leu Asp Val Leu Val Glu Met Gly					
	805			810		815
His Lys Glu Leu Lys Glu Ile Gly Ile Asn Ala Tyr Gly His Arg His						
	820			825		830
Lys Leu Ile Lys Gly Val Glu Arg Leu Ile Ser Gly Gln Gln Gly Leu						
	835			840		845
Asn Pro Tyr Leu Thr Leu Asn Thr Ser Gly Ser Gly Thr Ile Leu Ile						
	850			855		860
Asp Leu Ser Pro Asp Asp Lys Glu Phe Gln Ser Val Glu Glu Glu Met						
	865			870		875
Gln Ser Thr Val Arg Glu His Arg Asp Gly Gly His Ala Gly Gly Ile						
	885			890		895
Phe Asn Arg Tyr Asn Ile Leu Lys Ile Gln Lys Val Cys Asn Lys Lys						
	900			905		910
Leu Trp Glu Arg Tyr Thr His Arg Arg Lys Glu Val Ser Glu Glu Asn						
	915			920		925
His Asn His Ala Asn Glu Arg Met Leu Phe His Gly Ser Pro Phe Val						
	930			935		940
Asn Ala Ile Ile His Lys Gly Phe Asp Glu Arg His Ala Tyr Ile Gly						
	945			950		955
Gly Met Phe Gly Ala Gly Ile Tyr Phe Ala Glu Asn Ser Ser Lys Ser						
	965			970		975
Asn Gln Tyr Val Tyr Gly Ile Gly Gly Gly Thr Gly Cys Pro Val His						
	980			985		990
Lys Asp Arg Ser Cys Tyr Ile Cys His Arg Gln Leu Leu Phe Cys Arg						
	995			1000		1005
Val Thr Leu Gly Lys Ser Phe Leu Gln Phe Ser Ala Met Lys Met Ala						
	1010			1015		1020
His Ser Pro Pro Gly His His Ser Val Thr Gly Arg Pro Ser Val Asn						
	1025			1030		1035
Gly Leu Ala Leu Ala Glu Tyr Val Ile Tyr Arg Gly Glu Gln Ala Tyr						
	1045			1050		1055
Pro Glu Tyr Leu Ile Thr Tyr Gln Ile Met Arg Pro Glu Gly Met Val						
	1060			1065		1070
Asp Gly						

Claims:

1. An isolated polynucleotide molecule encoding a candidate effector protein for the Grb7 family of signalling proteins, wherein the polynucleotide molecule comprises a nucleotide sequence having at least 75% sequence identity to that shown as SEQ ID NO: 1.
2. A polynucleotide molecule according to claim 1, wherein the polynucleotide molecule comprises a nucleotide sequence having at least 85% sequence identity to that shown as SEQ ID NO: 1.
3. A polynucleotide molecule according to claim 1, wherein the polynucleotide molecule comprises a nucleotide sequence having at least 95% sequence identity to that shown as SEQ ID NO: 1.
4. A polynucleotide molecule according to claim 1, wherein the polynucleotide molecule comprises a nucleotide sequence which substantially corresponds to that shown as SEQ ID NO: 1.
5. A host cell transformed with a polynucleotide molecule according to any one of the preceding claims.
6. A host cell according to claim 5, wherein the host cell is a mammalian, insect, yeast or bacterial host cell.
7. A method of producing a protein, comprising culturing the host cell of claim 5 or 6 under conditions suitable for the expression of the polynucleotide molecule and optionally recovering the protein.
8. A purified protein encoded by a polynucleotide molecule according to any one of claims 1 to 4.
9. A purified protein according to claim 8, wherein the protein comprises an amino acid sequence substantially corresponding to that shown as SEQ ID NO: 2.

10. A fusion protein comprising an amino acid sequence substantially corresponding to that shown as SEQ ID NO: 2.
11. An antibody or fragment thereof which specifically binds to a protein
5 according to claim 8 or 9.
12. An oligonucleotide probe comprising a nucleotide sequence of at least 12 nucleotides, the oligonucleotide probe comprising a nucleotide sequence such that the oligonucleotide probe selectively hybridises to the
10 polynucleotide molecule of any one of claims 1 to 4 under high stringency conditions.
13. An oligonucleotide probe according to claim 12. wherein the oligonucleotide probe comprises a nucleotide sequence of at least 18
15 nucleotides.
14. A method of detecting in a sample the presence of an effector protein for the Grb7 family of proteins, the method comprising reacting the sample with an antibody or fragment thereof according to claim 11.
20
15. A method of detecting in a sample the presence of mRNA encoding an effector protein for the Grb7 family of proteins, the method comprising reacting the sample with an oligonucleotide probe of claim 12 or 13.

1/4

FIGURE 1

ATTCTCTTTCATAATGCATGCTCTTTTGGTCATGCTGAAGTAGTCAATCTCCTTTTGGCAGATGGTGCAG 70
I P L H N A C S F G H A E V V N L L L R H G A

ACCCCAATGCTCGAGATAATTGGAATTATACTCCTCTCCATGAAGCTGCAATTAAAGGAAAGATTGATGT 140
D P N A R D N W N Y T P L H E A A I K G K I D V

TTGCATTGTGCTGTACAGCATGGAGCTGAGCCAACCATCCGAAATACAGATGGAAGGACAGCATTGGAT 210
C I V L L Q H G A E P T I R N T D G R T A L D

TTAGCAGATCCATCTGCCAAAGCAGTGCTTACTGGTGAATATAAGAAAGATGAACCTTTAGAAAGTGCCA 280
L A D P S A K A V L T G E Y K K D E L L E S A

GGAGTGGCAATGAAGAAAAATGATGGCTCTACTCACACCATTAAATGTCAACTGCCACGCAAGTGATGG 350
R S G N E E K M M A L L T P L N V N C H A S D G

CAGAAAGTCAACTCCATTACATTTGGCAGCAGGATATAACAGAGTAAAGATTGTACAGCTGTTACTGCAA 420
R K S T P L H L A A G Y N R V K I V Q L L L Q

CATGGACGTGATGCCATGCTAAAGATAAAGGTGATCTGGTACCATTACACAATGCCTGTTCTTATGGTC 490
H G R D V H A K D K G D L V P L H N A C S Y G

ATTATGAAGTAACTGAACCTTTGGTCAAGCATGGTGGCTGTGTAATGCAATGGACTTGTGGCAATTCAC 560
H Y E V T E L L V K H G G C V N A M D L W Q F T

TCCTCTTCATGAGGCAGCTTCTAAGAACAGGGTTGAAGTATGTTCTCTTCTTAAAGTTATGGTGCAGAC 630
P L H E A A S K N R V E V C S L L L S Y G A D

CCAACACTGCTCAATTGTAAGAATAAAAGTGCTATAGACTTGGCTCCCACACCACAGTTAAAAGAAAGAT 700
P T L L N C K N K S A I D L A P T P Q L K E R

TAGCATATGAATTTAAAGGCCACTCGTTGCTGCAAGCTGCACGAGAAGCTGATGTTACTCGAATCAAAAA 770
L A Y E F K G H S L L Q A A R E A D V T R I K K

ACATCTCTCTCTGGAATGGTGAATTTCAAGCATCCTCAAACACATGAAACAGCATTGCATTGTGCTGCT 840
H L S L E M V N F K H P Q T H E T A L H C A A

GCATCTCCATATCCCAAAAGAAAGCAAATATGTGAACCTGTTGCTAAGAAAAGGAGCAAACATCAATGAAA 910
A S P Y P K R K Q I C E L L L R K G A N I N E

AGACTAAAGAATCTTGACTCCTCTGCACGTGGCATCTGAGAAAGCTCATAATGATGTTGTTGAAGTAGT 980
K T K E F L T P L H V A S E K A H N D V V E V V

GGTGAAACATGAAGCAAAGGTTAATGCTCTGGATAATCTTGGTCAGACTTCTCTACACAGAGCTGCATAT 1050
V K H E A K V N A L D N L G Q T S L H R A A Y

TGTGGTCATCTACAAACCTGCCGCCTACTCCTGAGCTATGGGTGTGATCCTAACATTATATCCCTTCAGG 1120
C G H L Q T C R L L L S Y G C D P N I I S L Q

GCTTTACTGCTTTACAGATGGGAAATGAAATGTACAGCAACTCCTCCAAGAGGTATCTCATTAGGTAA 1190
G F T A L Q M G N E N V Q Q L L Q E G I S L G N

TTAGAGGCAGACAGACAATTGCTGGAAGCTGCAAAGGCTGGAGATGTGAAACTGTAAAAAACTGTGT 1260
S E A D R Q L L E A A K A G D V E T V K K L C

ACTGTTCAAGAGTGTCAACTGCAGAGACATTGAAGGGCGTCAGTCTACACCACTTCATTTTGCAGCTGGGT 1330
T V Q S V N C R D I E G R Q S T P L H F A A G

ATAACAGAGTGTCCGTGGTGAATATCTGCTACAGCATGGAGCTGATGTGCATGCTAAAGATAAAGGAGG 1400
Y N R V S V V E Y L L Q H G A D V H A K D K G G

CCTTGATCTTTGCACAATGCATGTTCTTACGGACATTATGAAGTTGCAGAACTTCTTGTAAACATGGA 1470
L V P L H N A C S Y G H Y E V A E L L V K H G

GCAGTAGTTAATGTAGCTGATTTATGGAATTTACACCTTTACATGAAGCAGCAGCAAAAGGAAATATG 1540
A V V N V A D L W K F T P L H E A A A K G K Y

AAATTGCAAACTTCTGCTCCAGCATGGTGCAGACCCTACAAAAAAAACAGGGATGGAATACTCCTTT 1610
E I C K L L L Q H G A D P T K K N R D G N T P L

2/4

GGATCTTGTAAAGATGGAGATACAGATATTTCAAGATCTGCTTAGGGGAGATGCAGCTTTGCTAGATGCT 1680
D L V K D G D T D I Q D L L R G D A A L L D A

GCCAAGAAGGGTTGTTTAGCCAGAGTGAAGAAGTTGTCTTCTCCTGATAATGTAAATTGCCGCGATACCC 1750
A K K G C L A R V K K L S S P D N V N C R D T

AAGGCAGACATTCAACACCTTTACATTTAGCAGCTGGTTATAATAATTTAGAAGTTGCAGAGTATTTGTT 1820
Q G R H S T P L H L A A G Y N N L E V A E Y L L

ACAACACGGAGCTGATGTGAATGCCCAAGACAAAGGAGGACTTATTCCTTTACATAATGCAGCATCTTAC 1890
Q H G A D V N A Q D K G G L I P L H N A A S Y

GGGCATGTAGATGTAGCAGCTCTACTAATAAAGTATAATGCATCTCTCAATGCCACGGACAAATGGGCTT 1960
G H V D V A A L L I K Y N A S L N A T D K W A

TCACACCTTTGCACGAAGCAGCCCCAAAGGGACGAACACAGCTTTGTGCTTTGTTGCTAGCCCATGGAGC 2030
F T P L H E A A Q K G R T Q L C A L L L A H G A

TGACCCGACTCTTAAAAATCAGGAAGGACAAACACCTTTAGATTAGTTTTCAGCAGATGATGTGAGCGCT 2100
D P T L K N Q E G Q T P L D L V S A D D V S A

CTTCTGACAGCAGCCATGCCCCATCTGCTCTGCCCTCTGTTACAAGCCTCAAGTGCTCAATGGTGTA 2170
L L T A A M P P S A L P S C Y K P Q V L N G V

GAAGCCCAGGAGCCACTGCAGATGCTCTCTCTCAGGTCCATCTAGCCCATCAAGCCTTCTGCAGCCAG 2240
R S P G A T A D A L S S G P S S P S S L S A A S

CAGTCTTGACAACCTATCTGGGAGTTTTCAGAACTGTCTTCAGTAGTTAGTTCAAGTGGAACAGAGGGT 2310
S L D N L S G S F S E L S S V V S S S G T E G

GCTTCCAGTTTGGAGAAAAAGGAGGTTCCAGGAGTAGATTTTAGCATAACTCAATTCGTAAGGAATCTTG 2380
A S S L E K K E V P G V D F S I T Q F V R N L

GACTTGAGCACCTAATGGATATATTTGAGAGAGAACAGATCACTTTGGATGTATTAGTTGAGATGGGGCA 2450
G L E H L M D I F E R E Q I T L D V L V E M G H

CAAGGAGCTGAAGGAGATTGGAATCAATGCTTATGGACATAGGCACAACTAATTAAGGAGTCGAGAGA 2520
K E L K E I G I N A Y G H R H K L I K G V E R

CTTATCTCCGGACAACAAGGTCTTAACCCATATTTAACTTTGAACACCTCTGGTAGTGAACAATCTTA 2590
L I S G Q Q G L N P Y L T L N T S G S G T I L

TAGATCTGTCTCCTGATGATAAAGAGTTTCAGTCTGTGGAGGAAGAGATGCAAAGTACAGTTCGAGAGCA 2660
I D L S P D D K E F Q S V E E E M Q S T V R E H

CAGAGATGGAGGTCATGCAGGTGGAATCTTCAACAGATAACAATATTCTCAAGATTCAGAAGGTTTGTAA 2730
R D G G H A G G I F N R Y N I L K I Q K V C N

AAGAACTATGGGAAAGATACACTCACCGGAGAAAAAGAGTTTCTGAAGAAAACCACAACCATGCCAATG 2800
K K L W E R Y T H R R K E V S E E N H N H A N

AACGAATGCTATTTTCATGGGTCTCCTTTTGTGAATGCAATTATCCACAAAGGCTTTGATGAAAGGCATGC 2870
E R M L F H G S P F V N A I I H K G F D E R H A

GTACATAGGTGGTATGTTTGGAGCTGGCATTATTTTGTGCTGAAAACCTCTTCCAAAAGCAATCAATATGTA 2940
Y I G G M F G A G I Y F A E N S S K S N Q Y V

TATGGAATTGGAGGAGTACTGGGTGTCAGTTCACAAAGACAGATCTTGTACATTGCCCACAGGCAGC 3010
Y G I G G G T G C P V H K D R S C Y I C H R Q

TGCTCTTTTGGCGGTAACCTTGGGAAAGTCTTTCCTGCAGTTCAGTGCAATGAAAATGGCACATTCTCC 3080
L L F C R V T L G K S F L Q F S A M K M A H S P

TCCAGGTCATCACTCAGTCACTGGTAGGCCAGTGTAATGGCCTAGCATTAGCTGAATATGTTATTTAC 3150
P G H H S V T G R P S V N G L A L A E Y V I Y

AGAGGAGAACAGGCTTATCCTGAGTATTTAATTACTTACCAGATTATGAGGCCTGAAGGTATGGTCGATG 3220
R G E Q A Y P E Y L I T Y Q I M R P E G M V D

GATAAATAGTTATTTTAAGAACTAATTCCTGAACCTAAAATCATCAAAGCAGCAGTGGCCTCTACGT 3290
G *

3/4

TTTACTCCTTTGCTGAAAAAAAAATCATCTTGCCACAGGCCTGTGGCAAAGGATAAAAAATGTGAACGAA 3360

GTTTAACATTCTGACTTGATAAAGCTTTAATAATGTACAG

4/4

A**CONSTRUCT****STRUCTURE****N****C****N + C****FL****B****CONSTRUCT**
**MEAN RLU
(LIQUID ASSAY)
(X 10³)**
**COLOUR INTENSITY
(FILTER ASSAY)**
pAS2.1**4****-****N****109****++****C****3****-****N + C****194****++****FL****242****+++**

FIGURE 2

INTERNATIONAL SEARCH REPORT

 International application No.
PCT/AU 98/00795
A. CLASSIFICATION OF SUBJECT MATTER
 Int Cl⁶: C12N 15/11, 15/12; C07K 14/46, 19/00, 16/18; G01N 33/68; C12Q 1/68

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED
 Minimum documentation searched (classification system followed by classification symbols)
 See Electronic Databases

 Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched
 See Electronic Databases

 Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)
WPAT (DGENE) - SEQ.ID.NO:2; Genbank, EMBL, Swiss-prot, PIR - SEQ.ID.NO:1, SEQ.ID.NO:2;
MEDLINE - Grb7, Grb#, growth factor receptor bound
C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	Janes PW et al. (1997) "Structural determinants of the interaction between the erbB2 receptor and the Src homology 2 domain of Grb7". The Journal of Biological Chemistry volume 272(13) pages 8490-8497. See entire document	1-15
A	Keegen K and Cooper JA "Use of the two hybrid systems to detect the association of the protein-tyrosine-phosphatase, SHPTP2, with another SH2-containing protein, Grb7" Oncogene volume 12, pages 1537-1544. See entire document	1-15

☐ Further documents are listed in the continuation of Box C

☐ See patent family annex

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier application or patent but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

 Date of the actual completion of the international search
 4 November 1998

Date of mailing of the international search report

11 NOV 1998

 Name and mailing address of the ISA/AU
 AUSTRALIAN PATENT OFFICE
 PO BOX 200
 WODEN ACT 2606
 AUSTRALIA
 Facsimile No.: (02) 6285 3929

Authorized officer


JULIE CAIRNDUFF

Telephone No.: (02) 6283 2545